RADAR OBSERVATIONS OF MJO/WAVE INTERACTIONS DURING DYNAMO/CINDY2011/AMIE

Amanda DePasquale, Courtney Schumacher, and Anita Rapp

Department of Atmospheric Science, Texas A&M University

1. Introduction

Total Daily OLR During DYNAMO/CINDY2011/AMIE

The Madden-Julian Oscillation (MJO) initiates over the Indian Ocean and propagates into the western Pacific as a series of convective events. These convective events may interact with other convectively-coupled waves, like Kelvin and Equatorial Rossby waves. The DYNAMO/CINDY2011/AMIE field campaigns from October 2, 2011 to February 9, 2012, collected data to better understand MJO initiation. On Gan Island in the Maldives, Texas A&M University deployed a C-band Doppler radar, called SMART-R.

2. Event Identification

- Convectively coupled equatorial waves are identified by wavenumber-frequency filtering of outgoing longwave radiation (OLR) using 3 different filters and plotted by OLR anomalies at a -10 W/m^2 threshold

	Wheeler	Schreck	Dynamo
MJO	30-96 days	30-100 days	20-100 days
Kelvin Wave	2.5-30 days	2.5-17 days	2.5-20 days

MJO Events

-5°S to 5°N   -10°S to 10°N   -15°S to 15°N

- Each filtering method identifies the first two strong MJO events with less agreement in December and January and as the domain size increases

Kelvin Waves

-5°S to 5°N   -10°S to 10°N   -15°S to 15°N

- The Dynamo method captures a combination of the Wheeler and Schreck methods
- Narrow domains capture more Kelvin waves

3. SMART-R Rain Rate

Convective and Stratiform Rain Over Gan From SMART-R

Active MJO at Gan averaged from -10°S to 10°N

- Agreed on by all methods
- Wheeler
- Schreck
- Dynamo

4. Average Rain Rates by Domain

- Composite MJO and Kelvin wave rain rates and stratiform percentages calculated based on identification method and domain

Composite MJO

-5°S to 5°N   -10°S to 10°N   -15°S to 15°N

- MJO events always have more stratiform rain than convective rain
- Stratiform percent is consistent at different domain size for Wheeler and Dynamo methods, but increases with domain size for the Schreck method
- Total rain increases as event identification domain gets larger due to the reduction in events sampled

Composite Kelvin Wave

-5°S to 5°N   -10°S to 10°N   -15°S to 15°N

- Kelvin waves have more stratiform rain than convective rain until the largest domain for event identification
- All three methods are in good agreement using the narrowest domain for event identification
- As the domain widens, stratiform percent decreases and convective rain dominates, due to fewer identified events

5. Kelvin Wave Interaction

- Using the Dynamo method and averaging from -10°S to 10°N, 8 Kelvin waves and 3 MJO events were identified at Gan

SMART-R Rain Rates Centered at Kelvin Wave Passage

Suppressed MJO (11/9)   *Active MJO (11/28)*

- Suppressed MJO: convective rain increases to a maximum one day after the Kelvin wave passes
- Active MJO: convective rain spikes at the passage of the Kelvin wave, followed by a maximum of stratiform rain

Zonal Wind Anomalies Centered at Kelvin Wave Passage

Suppressed MJO (11/9)   *Active MJO (11/28)*

- Suppressed MJO: zonal winds from soundings show a vertical enhancement of easterly anomalies leading up to the wave passage followed by westerly anomalies after the wave passes over Gan
- Active MJO: zonal winds are dominated by MJO (strong upper-level easterlies and strong lower-level westerlies), but the vertical enhancement of easterly anomalies at the Kelvin wave passage is evident

6. Conclusions

- To study wave interactions, careful consideration of filtering method and domain size is necessary
- While smaller domain size is better for identification of shorter, faster Kelvin waves, larger domains are better for MJO events
- There is an enhancement of convective rain during a Kelvin wave passage, but significantly more stratiform rain during an active MJO, even though the easterly wind anomalies are similar
- More work is needed to examine the interaction of the MJO with Equatorial Rossby waves and consider other environmental properties like humidity

References: Wheeler M, and Kiladis GN. 1999. Convectively-coupled equatorial waves: Analysis of clouds in the wavenumber-frequency domain. *J. Atmos. Sci.* 56: 374-399.
Carl Schreck's website: http://monitor.cicsnc.org/mjo/current/